

## CORE CLINICAL QUALITY MEASURES

### Instructions:

1. EPs must complete 3 of the Core or Alternate Core CQM to satisfy MU requirements
2. Select each of the Core CQM that are being reported
3. If your EP's data is not available and this is a test run evaluation, select "data not available"
4. If your EP is reporting on Alternate Core, proceed to next grid

Core CQM #1	Numerator	Denominator	Data Not Available	Selected
Hypertension/ BP Measurement			<input type="checkbox"/>	<input type="checkbox"/>
Core CQM #2	Numerator	Denominator	Data Not Available	Selected
Tobacco Use Assessment			<input type="checkbox"/>	<input type="checkbox"/>
Tobacco Cessation Intervention			<input type="checkbox"/>	<input type="checkbox"/>
Core CQM #3	Numerator	Denominator	Data Not Available	Selected
Adult Weight Screening: Age 18-64			<input type="checkbox"/>	<input type="checkbox"/>
Adult Weight Screening: Age 65+			<input type="checkbox"/>	<input type="checkbox"/>

**ALTERNATE CORE MEASURES**

**Instructions:**

1. EP's must complete 3 of the Core or Alternate Core CQM to satisfy MU requirements
2. Select each of the Alternate Core that are being reported
3. If your EP's data is not available and this is a test run evaluation, select "data not available"
4. If your EP is reporting on Core CQM, use previous grid

Alt CQM #1	Numerator	Denominator	Data Not Available	Selected
BMI Percentile: Ages 2-16			<input type="checkbox"/>	<input type="checkbox"/>
Nutrition Counseling: Ages 2-16			<input type="checkbox"/>	<input type="checkbox"/>
Physical Activity Counseling: Ages 2-16			<input type="checkbox"/>	<input type="checkbox"/>
BMI Percentile: Ages 2-10			<input type="checkbox"/>	<input type="checkbox"/>
Nutrition Counseling: Ages 2-10			<input type="checkbox"/>	<input type="checkbox"/>
Physical Activity Counseling: Ages 2-10			<input type="checkbox"/>	<input type="checkbox"/>
BMI Percentile: Ages 11-16			<input type="checkbox"/>	<input type="checkbox"/>
Nutrition Counseling: Ages 11-16			<input type="checkbox"/>	<input type="checkbox"/>
Physical Activity Counseling: Ages 11-16				<input type="checkbox"/>
Alt CQM #2	Numerator	Denominator	Data Not Available	Selected
Influenza Immunization for patients >=50			<input type="checkbox"/>	<input type="checkbox"/>

<b>Alt CQM #3</b>	<b>Numerator</b>	<b>Denominator</b>	<b>Data Not Available</b>	<b>Selected</b>
<b>Childhood Immunization-DTaP</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization-IPV</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- MMR</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization-HiB</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- Hep B</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- VZV</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- PCV</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- HepA</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- RV</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- Flu</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- Combination Rate 1</b>			<input type="checkbox"/>	<input type="checkbox"/>
<b>Childhood Immunization- Combination Rate 2</b>			<input type="checkbox"/>	<input type="checkbox"/>

## ADDITIONAL CLINICAL QUALITY MEASURES

### Instructions:

1. Select 3 additional CQMs from the list below

Title	Description	Numerator	Denominator	Selected
Asthma Assessment	% of patients aged 5-40 w/ a dx of asthma and who have been seen for at least 2 office visits, who were evaluated during at least one visit within 12 mos. for the frequency (numeric) and nocturnal asthma symptoms			<input type="checkbox"/>
Appropriate testing for children with pharyngitis	% of children 2-18 yrs of age who were dx'd with pharyngitis, dispensed an antibiotic and rec'd a group A streptococcus test for the episode			<input type="checkbox"/>
Initiation and engagement of alcohol and other drug dependence treatment: (a) initiation (b) engagement	% of adolescent and adult patients with a new episode of alcohol and other drug (AOD) dependence who initiate tx through an inpt AOD admission, outpt visit, intensive outpt encounter or partial hospitalization within 14 days of the dx and who initiated tx and who had 2 or more additional services w/ an AOD dx within 30 days of the initiation visit.			<input type="checkbox"/>
Prenatal Care: Screening for HIV	% of patients, regardless of age, who gave birth during a 12-mo period who were screened for HIV infection during the first or second prenatal care visit			<input type="checkbox"/>
Controlling HBP	% of patients 18-85 yrs of age who had a dx of hypertension and whose BP was adequately controlled during the measurement year			<input type="checkbox"/>
Smoking and tobacco use cessation: Medical assistance: a. Advising smokers and tobacco users to quit, b. Discussing smoking and tobacco use cessation medications, c. Discussing smoking and tobacco use cessation strategies	% of patients 18 yrs of age and older who were current smokers or tobacco users, who were seen by a practitioner during the measurement year and who received advice to quit smoking or tobacco use or whose practitioner recommended or discussed smoking or tobacco use cessation medications, methods or strategies			<input type="checkbox"/>
Breast cancer screening	% of women 40-69 yrs of age who had a mammogram to screen for breast cancer			<input type="checkbox"/>
Cervical cancer screening	% of women 21-64 years of age, who received one or more pap tests to screen for cervical cancer			<input type="checkbox"/>
Chlamydia screening for women	% of women 15-24 yrs of age who were identified as sexually active and who had a			<input type="checkbox"/>

	least one test for Chlamydia during the measurement year			
Colorectal cancer screening	% of adults 50-75 yrs of age who had appropriate screening for colorectal cancer			<input type="checkbox"/>
Use of appropriate medications for asthma	% of patients 5-50 yrs of age who were identified as having persistent asthma and were appropriately prescribed medication during the measurement year. Report 3 age stratifications (5-11 yrs, 12-50 yrs, and total)			<input type="checkbox"/>
Pneumonia vaccination status for older adults	% of patients 65 yrs of age and older who have ever received a pneumonia vaccine			<input type="checkbox"/>
Asthma pharmacologic therapy	% of patients aged 5-40 yrs with a dx of mild, moderate, or severe persistent asthma who were prescribed either the preferred long-term control medication (inhaled corticosteroid) or an acceptable alt treatment			<input type="checkbox"/>
Low back pain: use of imaging studies	% of patients w/ a primary dx of low back pain who did not have an imaging study (plain x-ray, MRI, CT scan) within 28 days of dx			<input type="checkbox"/>
Diabetes: eye exam	% of pts 18-75 yrs of age w/ diabetes (type 1 or type 2) who had a retinal or dilated eye exam or a negative retinal exam (no evidence of retinopathy) by an eye care professional			<input type="checkbox"/>
Diabetes: foot exam	% of pts aged 18-75 yrs w/ diabetes (type 1 or type 2) who had a foot exam (visual inspection, sensory exam w/ monofilament, or pulse exam)			<input type="checkbox"/>
Diabetes: hemoglobin A1c poor control	% of pts 18-75 yrs w/ diabetes (type 1 or type 2) who had hemoglobin A1c > 9.0%			<input type="checkbox"/>
Diabetes: blood pressure management	% of pts 18-75 yrs w/ diabetes (type 1 or type 2) who had blood pressure <140/90 mmHg			<input type="checkbox"/>
Diabetes: urine screening	% of pts 18-75 yrs of age w/ diabetes (type 1 or type 2) who had a nephropathy screening test or evidence of nephropathy			<input type="checkbox"/>
Diabetes low density lipoprotein (LDL) mgmt & control	% of pts 18-75 yrs of age w/ diabetes (type 1 or type 2) who had LDL-C <100 mg/dL			<input type="checkbox"/>
Coronary artery disease (CAD): oral antiplatelet therapy prescribed for patients w/ CAD	% of pts aged 18 yrs and older w/ a dx of CAD who were prescribed oral antiplatelet therapy			<input type="checkbox"/>
Ischemic Vascular Disease (IVD: Use of aspirin or another antithrombotic	% of patients 18 yrs of age and older who were discharged alive for acute myocardial infarction (AMI), coronary bypass graft (CABG) or percutaneous transluminal coronary angioplasty (PTCA) from Jan 1-Nov			<input type="checkbox"/>

	1 of the year prior to the measurement year, or who had a dx of ischemic vascular dz (IVD) during the measurement year and the year prior to the measurement year and who had documentation of use of aspirin or another antithrombotic during the measurement year			
Coronary Artery Disease (CAD): Beta-Blocker therapy for CAD patients with prior myocardial infarction (MI)	% of patients 18 years of age and older with a dx of CAD and prior MI who were prescribed beta-blocker therapy.			<input type="checkbox"/>
Ischemic Vascular Disease (IVD): Blood pressure management	% of patients 18 yrs of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous transluminal coronary angioplasty (PTCA) from Jan 1-Nov 1 of the year prior to the measurement year, or who had a dx of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year and whose recent blood pressure is in control (<140/90mmHg)			<input type="checkbox"/>
Coronary Artery Disease (CAD): Drug therapy for lowering LDL-cholesterol	% of patients aged 18 yrs and older with a dx of CAD who were prescribed a lipid lowering therapy (based on current ACC/AHA guidelines)			<input type="checkbox"/>
Ischemic Vascular Disease (IVD): Complete lipid panel and LDL control	% of patients 18 yrs of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous transluminal angioplasty (PTCA) from Jan 1-Nov 1 of the year prior to the measurement year, or who had a dx of ischemic vascular dz (IVD) during the measurement year and the year prior to the measurement year and who had a complete lipid profile performed during the measurement year and whose LDL-C<100 mg/dL			<input type="checkbox"/>
Heart Failure (HF): Angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) therapy for left ventricular systolic dysfunction (LVSD)	% of patients aged 18 years and older with a dx of heart failure and LVSD (LVEF <40%) and who were prescribed ACE inhibitor or ARB therapy			<input type="checkbox"/>

<b>Heart Failure (HF): Beta-Blocker Therapy for left ventricular systolic dysfunction (LVSD)</b>	<b>% of pts aged 18 years and older with a dx of heart failure who also have LVSD (LVEF &lt;40%) and who were prescribed beta-blocker therapy</b>			<input type="checkbox"/>
<b>Heart Failure (HF): Warfarin therapy patients w/ atrial fibrillation warfarin therapy</b>	<b>% of all patients aged 18 yrs and older w/ a dx of heart failure and paroxysmal or chronic atrial fibrillation who were prescribed</b>			<input type="checkbox"/>
<b>Primary Open Angel Glaucoma (POAG): Optic Nerve Evaluation</b>	<b>% of patients aged 18 yrs and older w/ a dx of POAG who have been seen for at least two office visits who have an optic nerve head evaluation during one or more office visits within 12 mos</b>			<input type="checkbox"/>
<b>Diabetic retinopathy: documentation of presence or absence of macular edema and level of severity of retinopathy</b>	<b>% of patients aged 18 yrs and older w/ a dx of diabetic retinopathy who had dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy and the presence or absence of macular edema during one or more office visits within 12 mos</b>			<input type="checkbox"/>
<b>Diabetic retinopathy: communication with the physician managing ongoing diabetes care</b>	<b>% of patients aged 18 yrs and older w/ a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed and documented communication to the physician who manages the ongoing care of the patient w/ diabetes mellitus regarding the findings of the macular or fundus exam at least once within 12 mos</b>			<input type="checkbox"/>
<b>Anti-depressant medication management: (a) Effective Acute Phase Treatment, (b) Effective Continuation Phase Treatment</b>	<b>% of patients 18 yrs of age or older who were dx'd w/ a new episode of major depression, treated w/ antidepressant medication, and who remained on an antidepressant medication treatment</b>			<input type="checkbox"/>
<b>Oncology Colon Cancer: Chemotherapy for State III colon cancer patients</b>	<b>% of patients aged 18 yrs or older with Stage IIIA through IIIC colon cancer who are referred for adjuvant chemotherapy, prescribed adjuvant chemotherapy, or have previously rec'd adjuvant chemotherapy within the 12 month reporting period</b>			<input type="checkbox"/>

<b>Oncology Breast Cancer: Hormonal therapy for Stage IC-IIIC estrogen receptor/progesterone receptor (ER/PR) positive breast cancer</b>	<b>% of female patients aged 18 yrs and older w/ Stage IC through IIIC, ER or PR positive breast cancer who were prescribed tamoxifen or aromatase inhibitor (A1) during the 12 month reporting period</b>			<input type="checkbox"/>
<b>Prostate Cancer: Avoidance of overuse of bone scan for staging low risk prostate cancer patients</b>	<b>% of patients, regardless of age, w/ a dx of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since dx of prostate cancer</b>			<input type="checkbox"/>
<b>Diabetes: Hemoglobin A1c Control (&lt;8.0%)</b>	<b>The percentage of patients 18-75 yrs of age w/ diabetes (type 1 or type 2) who had hemoglobin A1c &lt;8.0%</b>			<input type="checkbox"/>